Online at: https://researchtrendsjournal.com **ISSN No:** 2584-282X
Indexed Journal **Peer Reviewed Journal**

INTERNATIONAL JOURNAL OF TRENDS IN EMERGING RESEARCH AND DEVELOPMENT

Volume 3; Issue 6; 2025; Page No. 37-42

Received: 14-08-2025
Accepted: 22-10-2025
Published: 16-11-2025

Applications of Group Actions in Algebraic Structures

[1]Veena Harode and [2]Pratima Ojha

[1, 2]Department of Mathematics, Madhyanchal Professional University, Bhopal, Madhya Pradesh, India

DOI: https://doi.org/10.5281/zenodo.17809973

Corresponding Author: Veena Harode

Abstract

In abstract algebra and other branches of mathematics, the concept of a large group of individuals collaborating on a gives a strong foundation for comprehending how a group can symmetrically transform the elements of a set. The thesis centers on exploring and resolving the complexities associated with group actions on sets, particularly in terms of their advanced theoretical aspects and practical applications. This research seeks to push the boundaries of current knowledge, providing new insights and tools for both theoretical and applied mathematics. Using software tools like Mathematica, MATLAB, or GAP (Groups, Algorithms, and Programming) to simulate group actions and visualize their effects on sets and to implement algorithms for solving group action-related problems, such as computing orbits, stabilizers, and automorphism groups. This study helps in analyzing the efficiency and scalability of these algorithms, particularly in handling large or complex groups.

Keywords: Applications, Group Actions, Algebraic Structures, MATLAB, GAP

Introduction

In abstract algebra, Group action theory offers a robust basis for understanding how algebraic structures interact with sets. A group action describes a way in inside a group, which includes a collection of components and a process that satisfies certain axioms, systematically "acts" on a set, thereby providing insight into its structure and symmetries. This concept extends beyond pure algebra into geometry, combinatorics, and even theoretical physics, making it an essential topic in higher mathematics.

At its core, A group action is a function that links each element of the group with a set while associating with maintaining the group structure. The presence of a function $G \times X \rightarrow XG$ would provide a more exact description of an action of a group G on a set X to $XG \times X \rightarrow X$. satisfies two fundamental properties: identity preservation (X undergoes an identity transformation via the part of G that is uniquely identified, and the process guarantees compatibility of a product of two group elements is the same as applying them successively). These conditions ensure that the group elements induce well-behaved transformations on the set, leading to significant structural insights.

The study of group actions is crucial because it connects abstract algebraic groups with more tangible geometric and combinatorial objects. One of the fundamental uses of group actions is in classifying objects up to symmetry. For example, in geometry, group actions help describe the symmetries of shapes, such as the rotational and reflectional symmetries of a square or a tetrahedron. In combinatorics, they provide a systematic way to count objects while considering symmetries, as seen in Burnside's Lemma, which aids in counting distinct objects under symmetry constraints.

Moreover, the study of group actions leads leading to significant findings like the orbit-stabilizer theorem, which describes how elements of a set are partitioned into orbits under a group action and relates this to subgroup properties. The concept of orbits (the set of points a particular element can move to under group actions) and stabilizers (subgroups of elements that leave a point unchanged) are fundamental in understanding how algebraic structures impose order and classification in mathematical spaces.

Beyond pure mathematics, group actions have applications in various scientific fields. In physics, they play a key role in quantum mechanics and crystallography, where the symmetries of molecules and crystal lattices can be described using group actions. In computer science, they appear in coding theory and cryptography, particularly in algorithms related to permutations and transformations.

Literature Review

Ishita Goluguri and Christina Li (2019) [1] This study delves into some intriguing uses of group actions, a miniature version of the instruments used to examine symmetry in group theory. The first step in doing this is to learn the basics of group theory so that we can analyze and evaluate group behaviors. Groups and their actions are defined and shown using basic examples, such as the group of square symmetries and its action on a vertex. Next, we show the Orbit-Stabilizer Theorem, a key result in proving Burnside's Lemma, and we define an orbit and a stabilizer. Afterwards, we show how Burnside's Lemma applies to combinatorial issues. Specifically, using Burnside's Lemma, we determine how many different ways a geometric design may be colored and how many different necklaces can be created using colored beads. Lastly, we go over the Sylow Theorems, another use of group actions, that are fundamental for group classification. Among other applicable definitions, we demonstrate these theorems using the conjugation group action.

Daniel Guan (2023) [2] This post provides an update on my prior survey from about 2020, focusing on developments in the applications of group actions in complicated geometry that mostly pertain to my own interests.

Dinesh Khattar (2023) [3] Our discussion is based on collective action and is supported by several examples. One powerful method for understanding group dynamics is collective action. It is a way to show how a set's symmetries reflect the group's elements. As an example, as a natural group, the dihedral group D4 operates on square vertices when shown as a collection of square symmetries action for many groups. An extension of this concept that may be used to extract valuable information about the group and the set it operates on is the collective behaviour when applied to a set.

Jesse Elliott (2020) [4] One of our services is a method for studying musical scales using the idea of group activities. For each group G, finite G-set S, and real number t, the {\it t-power diameter} diamt(G, S) is the size of any maximal orbit of S divided by the t-power mean orbit size of the elements of S. If the set of all tonic scales is subject to the symmetric groups, then {\its tonic scale} is a subset of Z12 that includes 0 process Chapter 11. Every single G-set heptatonic scale has the highest t-power diameter provided by {(1 2), (3 4), (5 6), (8 9), (10 11)}, for every t ∈ [−1,1], for the subgroup œ and its conjugate subgroups. For any subgroup G of S11, this is true. Bhatkhande asserts that the 32 thās that comprise Hindustani classical music create an exceptional highest oe-orbit. Among the 462 heptatonic scales, this research explains why 32 of them are interesting from a mathematical perspective. Our approach is also somewhat extended to hexatonic and pentatonic scales.

Evelyn Zhu (2022) [5] We study typical theories, examples, and applications of symmetric groups in a symmetric group in this work. The group operation is function composition, and all elements are bijections to the set itself. We start by going over what symmetry is and how it works, then we'll go over several instances. Next, we will go over the characteristics and unique components of symmetric groups before introducing and defining a few practical applications. The subgroup structure of symmetric groups is then shown, along with a few illustrative hypotheses.

Research Methodology

Construct models that illustrate how group actions operate on sets in specific contexts, such as symmetries in geometry or permutations in combinatorics. Apply the orbit-stabilizer theorem and other relevant theorems to analyze the behavior of these models. Develop new theorems or generalizations, if possible, to extend the application of group actions to previously unexplored areas.

Utilize software tools like Mathematica, MATLAB, or GAP (Groups, Algorithms, and Programming) to simulate group actions and visualize their effects on sets. Implement algorithms for solving group action-related problems, such as computing orbits, stabilizers, and automorphism groups. Analyze the efficiency and scalability of these algorithms, particularly in handling large or complex groups.

Data Analysis

Group Action" Definition

The unit's central idea, the significance of "group action on a set," is introduced in this section. Some instances of collective acts are provided in the next section. In order to understand how symmetries contribute to the development of §4.1-(1) discusses the idea of collective activities.

Sets of G-radii: Assume that G is a group and X is a set. The statement "G acts on X" or "X is a G-set" may be used to describe a function G × X → X that meets certain axioms. The conjugate image of (g,x) is represented as g · x in this function (for the g axiom), where x is an element of X and g is an element of G:

1. $e \cdot x = x$ where x belongs to and e is the identity element of the group X G
2. $g \cdot (h \cdot x) = (gh) \cdot x$ gh is the product of g and h in G, in that order, for all x in X, every g in G, and all h in G.

Thoughts on symbols and language

You have X as a G-set and G as a group. The action is defined by the function G × X → X, which is referred to as the "action map". It is frequently easier to write gx instead of g · x, which depicts the image underlying the action map of a pair (g, x), to avoid the "dot" in that notation. This "abuse of notation" is supported by postulate (2) of §2.3-3.1 becomes $g(hx) = (gh)x$, It seems to be nothing more than the "associativity axiom" for the group's multiplication operation. Indeed, we will need to deal with actions when the set x is the group G itself. (see §4.3). In a few of them, the group multiplication serves as the action map (§4.3.1) that is, its "direct descendant" (e.g., §4-(c)). By replacing the dot with gx, of $g \cdot x$ would simplify and clarify the notation in these instances. But keep in mind that when you write gx for $g \cdot x$ may in It is best to avoid using it in some instances because of the potential for ambiguity and confusion caused by conflicts of notation. With respect to the "conjugation action" §4-(b), As an example, the group is once again the set that is acted upon, but this time $g \cdot x$ describes what gxg^{-1}(the product of x and g. A catastrophe would ensue to mitigate the dot in this case. Various forms of notation, including as g_x to denote gxg^{-1}, is able to both convey message and be aesthetically pleasing.

New way of looking at G-sets

Subsection 1 offers a different way of putting the phrase that is instructive and useful. Why don't we start moving in the right direction? For the sake of argument, let's pretend that G acts upon the set X. There is a function φ_g (X) → X that is defined for every member g of by x → g · x. G, $\varphi_g(x) := g \cdot x.$ Now we may state the two axioms of the definition in §3.1 in an analogous manner:

1. φ_e presents itself as an itinerary map X, where e represents the unique identifier for G.
2. $\varphi_g \circ \varphi_h = \varphi_{gh}$, where $\varphi_g \circ \varphi_h$ indicates that the combination (φ_h initially, and then φ_g)
3. It is highly suggested by the second axiom in its equivalent form (2'), which states that the connection $g \to \varphi_g$ is homomorphic to the group! To make this assumption more concrete, notice Therefore when (1') and (2') are taken together, they show that φ_g serves as a self-contained transitive closure of X, $\varphi_{g_{-1}}$ opposite of φ_g.

Now consider the group $\mathfrak{S}_x$ for every bijection involving X on its own. Note that $\mathfrak{S}_x$ is a group that is currently being composed, with the identification map on X serving as its identity element. This particular instance is probably not new to the reader: In cases as an example, consider the set of all positive integers as X, which is a finite set of n items from 1 to n, then it transforms into $\mathfrak{S}_x$ is denoted $\mathfrak{S}_n$ and went by the name of "symmetric group on n letters".

It the statement made by (2') is readily verifiable $\varphi : G \to \mathfrak{S}_x$ given by $g \to \varphi_g$ is a group homomorphism. On the other hand, when thought of as a group homomorphism $\varphi : G \to \mathfrak{S}_x$ due to an association between It is possible to get an action of G by applying the definition to the composition of group G and the bijections of set X.$g \cdot x = \varphi_g(x);$ this place, we have indicated by φ_g $(rather\ than\ \varphi(g))$ the image of g under φ to avoid the ugliness of $\varphi(g)(x)$.

a) A synopsis of this part. Seen as a group homomorphism from G to X, it's the same as acting on X from group G. $\mathfrak{S}_x$ X bijections examined. Given that for any g in G, we operate G on X and thereafter allow φ_g illuminate the way $x \to g \cdot x$, then φ_g connects two sets of data and the relationship $g \to \varphi_g$ proves that two groups are identical G to $\mathfrak{S}_x$. The other one hand, Group homomorphism exists $\varphi : G \to \mathfrak{S}_x$, setting $g \cdot x := \varphi_g(x)$, in which ϕg portrays an image of g in $\mathfrak{S}_x$ under φ, $G's$ X's influence is proven. Connecting a homomorphism to an action, and vice versa, is obviously inversely related.

G-maps between G-sets: So, let's pretend given that G contains the sets X and Y How Is it possible to determine whether X and Y, two variables, are "the same"? Remember when there were two sets are isomorphic; we treat them as one and the same. Does the concept of "sameness" hold for G-sets in a same way? We begin by defining a G-map so that we may proceed to present such a concept. A vector (or procedure) $\varphi : X \to Y$ Every map from one set of G-values where G-equivariant means that it is a G-map if $g(\varphi(x)) = \varphi(gx)$ if we are given x from X and g from G. We may say that φ is a G-map if it commutes with the action of G. An additional set of G-maps is itself another set of G-maps; this is readily verifiable. Bijective G-maps are one kind of G-isomorphism. A G-isomorphism is required for X to be isomorphic to Y as a G-set. Otherwise, X cannot be isomorphic to Y. (G) homomorphism $\varphi: X \to Y$, its polar opposite $\varphi^{-1} : Y \to X$ includes an authentic Google Map. On G-sets, the equivalence relation "X is isomorphic to Y" is specified. For any two isomorphic G-sets, we say that they are identical. Because there is a G-isomorphism between them, any part of G's activity on one of them is reflected in G's action on the other, therefore it makes sense.

Group Actions in Algebraic Structures

Groups and locations have a tight connection. There are several communities associated with a certain place or region. We may investigate the group of symmetries, which is also called the group of structure-preserving bijections. In addition to the homology and cohomology groups, to name a few more, there is the fundamental group. According to Hermann Weyl, these groups might provide "a deep insight" into a particular place. An excellent example of this phenomenon may be found in the study of knots. For instance, algebraic invariants that take the form of groups demonstrate that the trefoil knot cannot be unravelled (see figure 1).



Fig 1: Groups show that these knots are distinct

The field of geometric group theory takes a fresh look at the connection between spaces and groups. The basic idea of geometric group theory is to adopt the following philosophy rather than analyzing spaces using the algebraic structure and properties of groups. Using your results, explore the geometry and topology of the spaces the groups work on. In other words, groups are the main subjects of study, and the techniques, instruments, and procedures used to investigate them are dynamical, geometrical, and topological by definition. Compared to the titles of other branches of mathematics, the word "geometric group theory" is relatively new. The notion of hyperbolic groups and the study of finitely generated groups as metric spaces were first presented by Gromov in his major works [Gro87, Gro93], which sparked a great deal of research and established areas of investigation that are still very active today. This substantial quantity of study and the development of various avenues of investigation are attributed to Gromov's efforts. Geometric notions were previously included into group theory prior to the creation of geometric group theory. The works of writers like Dehn, Whitehead, van Kampen, and others include these ideas. Furthermore, Thurston's work on 3-manifolds showed how a manifold's geometry may influence the algebraic and computational properties of its fundamental group. However, Gromov's publications are the ones that mark the beginning of these ideas' prominence. The aim of this article is to provide some light on how the algebraic structure of groups operating on a space is affected by the topology and geometry of that space, and how this knowledge may be used to study group behaviour. I use the strategy that my mentor, Mladen Best vina, taught me, which is to give more weight to illuminating cases than general theory. You'll notice that I use this approach. As is the case with every review of a mathematical field, many aspects and areas of geometric group theory are not covered at all in this overview. The last section offers a selection of publications that serve as a concise bibliography of further reading material on geometric group theory.

Groups and spaces: Geometric group theory, as was just said, uses group actions on spaces to try to understand the structure of a group. What insights might one possibly get from seeing how something is done? Will there always be anything interesting to investigate? After capturing your interest, let's examine each of these enquiries.

An example $SL(2\mathbb{Z})$: To illustrate how the topology of the space a group acts on may affect the group's structure, let's look at an example of a group action that appears in a wide variety of mathematical subfields. Here, we will concentrate on the set of matrices that have integer entries and determinants equal to one. The members of the so-called special linear group are as follows:

$$SL(2,\mathbb{Z}) = \left\{ \begin{bmatrix} a & b \\ c & d \end{bmatrix} \mid a,b,c,d \in \mathbb{Z} \ \ and \ \ ad-bc=1 \right\}$$

Is $SL(2,Z)$ finitely generated? That is, are there finitely many matrices $A1,\ldots,An \in SL(2,Z)$ such that any matrix $M \in SL(2,Z)$ can be expressed as a product $M = A \pm 1j1 \ldots A \pm 1jk$? (Keep in mind that any Aj could show up more than once. The answer is "yes," and this topic may be approached algebraically. However, let's adopt a geometric viewpoint and examine an action of $SL(2\mathbb{Z})$ on a metric space

The Farey complex, which is built as follows, is the area we will be looking at. We begin with a graph whose vertex set consists of the set of rational numbers p q, which are always written in lowest terms, plus one extra point that we designate as 1 0. Edges join two vertices $p\ q$ and $r\ s$ if $ps - qr = \pm 1$. A section of this graph, referred to as the Farey graph, is seen in Figure 2. The Farey graph's edges naturally form triangles, as seen in Figure 2. In actuality, every such triangle's vertices always have the form $p\ q$, $r\ s$ and $p + rq + s$. For instance, 1 0, 0 1 and 1 1, and also 1 0, 1 1 and 2 1. There is an action of $SL(2\mathbb{Z})$ on the Farey graph that is created by using the rule to permute the vertices.

$$\begin{bmatrix} a & b \\ c & d \end{bmatrix} \cdot \frac{p}{q} = \frac{ap+bq}{cp+dq}$$

Verifying that two vertices is simple $\frac{p}{q}$ and $\frac{r}{s}$ are only joined by an edge if their pictures $\begin{bmatrix} a & b \\ c & d \end{bmatrix} \cdot \frac{p}{q}$



Fig 2: The Farey graph and Farey complex

And $\begin{bmatrix} a & b \\ c & d \end{bmatrix} \cdot \frac{r}{s}$ are. As a result, an action is specified on the Farey complex and, therefore, on the Farey graph. The space that results from filling in the triangles in the Farey graph is known as the Farey complex. You have undoubtedly already seen this place and the activities that go along with it, although in a different form. In actuality, the Farey complex uses ideal triangles, whose vertices are either rational or in the model of the upper half plane, to produce a tessellation of the hyperbolic plane. Furthermore, the previously stated action requires fractional linear transformations of the top half plane and is just the usual action of two-by-two matrices with real entries and a positive determinant. The following are the conformal maps:

$$f(z) = \frac{1 - iz}{z - i} \quad and \quad g(z) = \frac{1 + iz}{z + i}$$

The time to look into this activity is right now. We will use the symbol to identify the triangle in the Farey complex that has the vertices 1 0 and 0 1 as well as the triangle in the Farey complex that has the vertices 1 0 and 1 1. The most significant aspects of the lawsuit are captured by two claims.



Fig 3: The tessellation of the upper half plane by ideal triangles according to the Farey algorithm

Claim 1: For any triangle $\Delta 0$ in the Farey complex, there is matrix $M \in SL(2, Z)$ such that $M\Delta = \Delta 0$.

Indeed, suppose the vertices of Δ' are $\frac{p}{q}$ $\frac{r}{s}$ and $\frac{p+q}{r+s}$ where, take $M = [\begin{smallmatrix} p & q \\ r & s \end{smallmatrix}]$ and observe that $M\Delta = \Delta 0$.

Alternative procedures on spaces that are δ-hyperbolic: Quite a few categories in nature simply cannot hyperbolic because their subgroups are isomorphic to Z 2. Is it feasible for us to use negative curvature to further investigate these groups? As we loosen the need of a valid metric space and the restriction of a geometric action, we will consider illustrates a noteworthy group within the domain of topology with few dimensions. An orientation able surface, which may or may not have a border, and the MCG () of orientation-preserving homeomorphisms of modulo isotopy constitute a class group. Just to clarify, homeomorphisms that may be continuously deformed into each other such that each intermediate map is likewise a homeomorphism are considered to define the same mapping class. It is required that the homeomorphisms and the isotopies on each component of the boundary have the same identity when the boundary is not empty. Let's suppose that is compact to make this discussion easier to handle. This group is used in the field that studies symplectic geometry, algebraic geometry, dynamics, configuration spaces, and three-manifolds, as well as in cryptography. Using homeomorphisms provided on disjoint subsurface Discovering isomorphic subgroups to Z 2 is a breeze in the majority of mapping class groups. Consequently, the MCG () function is generally not hyperbolic. The graph of the mapping class group influences curve C (). The circle S 1 embedded on a plane without an annulus (which appears only when it has a boundary) and a disc is called a closed curve that is simple. The curve graph is the collection of basic closed curve isotopy classes that make up the vertex set of the graph. If the representations of two such curves, [c0] and [c1], do not have a discontinuous connection with one other, then they are connected by an edge. Numerous curves are shown in Figure 4; each accompanied with the corresponding subgraph of C (). The equation illustrates how Any time a mapping class [f] is applied to a vertex [c] in the curve graph, the basic closed curve is transferred to the vertex's image. This holds for an operation on C as well, as homeomorphisms convert disjoint curves into other disjoint curves.

When the genus is equal to 1, any two non-isotopic simple closed curves will always intersect, because does in the case of a torus S_ (1). The aforementioned definition will thus result in a graph devoid of edges. The concept is significantly altered in this case such that if [c0] and [c1] have representatives that overlap once, they are regarded as being connected by an edge. Let's examine this curve graph in further detail. A simple closed curve on the torus in its most basic form is isotopic to a curve that wraps p times around the first S () component.



Fig 4: The graph of a genus 2 surface's curves

where p and q are approximately prime, and for iterations q that concentrate on the second S 1 component and beyond. Given that q is probably positive, we may safely infer that it is it doesn't matter which way the vector is orientated. In other words, parameters for the set of rational integers p and q plus one additional element, 1.0, constitute the isotopy classes of simple closed curves on the torus. Also, how often do the fundamental closed curves p, q, r, and s occur, appear intersect is shown by the equation "ps qr." Does anybody recognise this? The graph of the torus's curve is the Farey graph, as you rightly pointed out. The two operations are equivalent simply because SL (2, Z) is genuinely isomorphic to the torus's mapping class group. An excellent example of the impact of SL (2, Z) on the Farey graph exemplifies many key features of C () and the impact of MCG () on C (). First, we discovered that, like the Farey graph, the curve graph has a hyperbolic form. Masur and Minsky [MM99] were the first to demonstrate this startling fact, although it has since been repeatedly refuted. The most exact estimate of when they proved that was less than 17 [HPW15], which was given by Hennsel, Przytycki, and Webb. The importance of this result for the study Geometric group theory, 3-manifold geometry, and the mapping class group are all very important. Secondly, the action's discontinuity is not well implemented. In actuality, the vertex stabilizers have no upper limit. But this is a strength rather than a weakness! When homeomorphisms fix a simple closed curve, they do precisely what they sound like.

Conclusion

Research into group actions bridges multiple areas of mathematics, offering solutions to problems in abstract algebra, geometry, combinatorics, and applied sciences. By understanding these concepts, mathematicians and scientists can classify objects more efficiently, simplify complex counting problems, and discover deep structural properties of mathematical and physical systems. The versatility and applicability of group actions make them one of the most powerful tools in modern mathematics and theoretical sciences.

References

1. Goluguri RR, Sen S, Udgaonkar J. Microsecond sub-domain motions and the folding and misfolding of the mouse prion protein. Elife. 2019;8:e44766.
2. Guan D, Liang MX. Complete co-homogeneity one Kähler metrics on the affine quadric of complex dimension two (related to a co-homogeneity one point of view on a Yau Conjecture) [J]. Accepted and to appear in Chinese Quart. J. Math. 2023.
3. Khattar D, Agrawal N, Sirohi M. Dynamical analysis of a 5D novel system based on Lorenz system and its hybrid function projective synchronisation using adaptive control. Pramana. 2023;97(2):76.
4. Elliott J, Bai Z, Hsieh SC, Kelly SE, Chen L, Skidmore B, *et al*. ALK inhibitors for non-small cell lung cancer: A systematic review and network meta-analysis. PloS one. 2020;15(2):e0229179.
5. Zhu F, Yan Y, Doyle E, Zhu C, Jin X, Chen Z, *et al*. Microplastics altered soil microbiome and nitrogen cycling: the role of phthalate plasticizer. Journal of Hazardous Materials. 2022;427:127944.
6. Arvind N, Panja S. On $\mathbb{Z}_n \rtimes \mathbb{Z}_2$-Hopf–Galois structures. Journal of Algebra. 2022;596:37–52.
7. Arvind N, Panja S. Unit group of $\mathbb{F}_q SL(3,2), p \geq 11$. arXiv:2106.07261.
8. Arvind N, Panja S. Unit group of some finite semisimple group algebras. Journal of the Egyptian Mathematical Society. 2022. doi:10.1186/s42787-022-00151-0.
9. Arvind N, Panja S. Hopf–Galois realizability of $\mathbb{Z}_{n^2}$. Journal of Pure and Applied Algebra. 2023;227(4).
10. Byott NP. Uniqueness of Hopf–Galois structure of separable field extensions. Communications in Algebra. 1996;24:3217–3228.
11. Bidwell JNS. Automorphisms of direct products of finite groups: II. Archiv der Mathematik (Basel). 2008;91(2):111–121.

Creative Commons (CC) License
This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY 4.0) license. This license permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.